

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

## Detection of Epstein-Barr virus DNA in cardiac and aortic tissues from chronic, active Epstein-Barr virus infection associated with Kawasaki disease-like coronary artery aneurysms

Hideaki Kikuta, MD, Yukio Sakiyama, MD, Shuzo Matsumoto, MD, Isamu Hamada, MD, Makoto Yazaki, MD, Toshimitsu Iwaki, MD, and Masao Nakano, MD

From the Department of Pediatrics, Hokkaido University School of Medicine, Sapporo, the Pediatric Clinic, National Hakodate Hospital, Hakodate, the Department of Pediatrics, Nagoya City University Medical School, Nagoya, and the Department of Pediatrics, Gifu Prefectural Tajimi Hospital, Gifu, Japan

**We describe three patients with chronic, active Epstein-Barr virus infection associated with Kawasaki disease-like coronary artery aneurysms. The Epstein-Barr virus genome was detected in three cardiac tissue samples and one aortic tissue sample examined by means of the polymerase chain reaction. These findings suggest that chronic Epstein-Barr virus infection may play a pathogenic role in the development of coronary artery aneurysms. (J PEDIATR 1993;123:90-2)**

Chronic, active Epstein-Barr virus infection is characterized by fever, lymphadenopathy, splenomegaly, hepatitis, interstitial pneumonitis, and uveitis. Laboratory findings include anemia, thrombocytopenia, leukopenia, and hypergammaglobulinemia with a very high titer of EBV antibodies<sup>1,2</sup>; cardiac involvement is rare. To our knowledge, our previous report of CEBV associated with coronary artery aneurysms was the first.<sup>3</sup> We now report three patients with CEBV associated with coronary artery aneurysms, similar to the findings of Kawasaki disease.<sup>4</sup>

### METHODS

**Subjects.** Three Japanese patients with CEBV and coronary artery aneurysms were studied (Table). Coronary artery aneurysms were detected by cross-section echocardiography. All patients had a chronic illness characterized by prolonged fever, coronary artery aneurysms, hepatosplenomegaly, liver dysfunction, and hypergammaglobulinemia. In addition to coronary artery aneurysms, patients 1 and 3 had dilation of the sinus of Valsalva, and patient 1 died of rupture of the sinus of Valsalva. Patients 2 and 3 also had pericarditis. These patients had several other manifestations of EBV infection: interstitial pneumonitis in patient

1, interstitial nephritis<sup>5</sup> in patient 2, and rash in patient 3 (Table). None had mucous membrane involvement. The DNA was extracted from different sites in three patients with CEBV and two control cardiac tissues. Human DNA samples from Raji cells and Molt-4 cells were used as positive and negative control samples, respectively. Human embryonic fibroblast cells infected with herpes simplex virus type 1 (Seibert strain), varicella-zoster virus (wild type), and human cytomegalovirus (wild type) and human herpes

CEBV	Chronic, active Epstein-Barr virus infection
EBV	Epstein-Barr virus
KD	Kawasaki disease
PCR	Polymerase chain reaction

virus-6-infected cord blood mononuclear cells were used for these studies.


**EBV serologic tests.** Antibody titers to the EBV viral capsid antigen and early antigen were determined by indirect immunofluorescence. Antibody titers to EBV-determined nuclear antigen were determined by anticomplement immunofluorescence.<sup>3</sup>

**Polymerase chain reaction.** The PCR was performed according to a previously described method.<sup>6</sup> The PCR reaction mixture consisted of 200  $\mu$ mol of each deoxyribonucleotide, 2.5 U of *Taq* DNA polymerase, 50 nmol of potassium chloride per liter, 10 mmol of Tris-HCl (pH 8.3) per liter, 1.5 mmol of magnesium chloride per liter, 0.01% (wt/vol) of gelatin, 20 pmol of each oligonucleotide primer, and 1  $\mu$ g of DNA in a volume of 100  $\mu$ l. Samples were then subjected

Submitted for publication Sept. 29, 1992; accepted Jan. 26, 1993.  
Reprint requests: Hideaki Kikuta, MD, Department of Pediatrics, Hokkaido University School of Medicine, N-15, W-7, Kita-ku, Sapporo 060, Japan.

Copyright © 1993 by Mosby-Year Book, Inc.  
0022-3476/93/\$1.00 + .10 9/22/45927

NOTICE: This material may be protected  
by copyright law (Title 17 US Code)



**Figure.** Ethidium bromide staining of PCR products. Aliquots (10  $\mu$ l) of PCR products were subjected to electrophoresis. Lane 1, Aortic tissue DNA from patient 2; lane 2, cardiac tissue DNA from patient 2; lane 3, cardiac tissue DNA from patient 3; lane 4, cardiac tissue DNA from patient 1; +, Raji cell DNA, positive control sample; -, Molt-4 cell DNA, negative control sample; bp, base pairs.

to 35 cycles of PCR, each consisting of 1 minute of denaturation at 94° C, 2 minutes of annealing at 55° C, and 3 minutes of polymerization at 72° C by a DNA thermal cycler (Perkin-Elmer Cetus, East Norwalk, Conn.). The PCR procedure was performed with primers 5'-GTGACTTCACCAAAGGTCAG-3' and 5'-TTAAAGTCCACTTACCTCTG-3'. The primers amplify and detect a 410-base-pair sequence in the region within the EBV *Bam*HI-W fragment.

**Southern blot hybridization.** Electrophoresis of 10  $\mu$ l of the PCR product was performed on 1.8% agarose gel. The gel was stained with 1  $\mu$ g of ethidium bromide per milliliter for 5 minutes, and the DNA bands were visualized under ultraviolet light. The PCR product was transferred onto nitrocellulose membrane filters. Each filter was hybridized with <sup>32</sup>P-labeled cloned fragment *Bam*HI-W of the EBV genome for 48 hours at 41° C in 1× standard saline citrate (0.15 mol sodium chloride per liter and 0.015 mol sodium citrate per liter), 50% formamide, 0.5% sodium dodecylsulfate, and heat-denatured salmon sperm DNA, 100  $\mu$ g/ml. After hybridization, the filters were washed three times at room temperature in 0.1× standard saline citrate with 0.1% sodium dodecylsulfate and then incubated three times for 1 hour at 50° C. The filters were dried and exposed to x-ray film at -80° C.<sup>7</sup>

## RESULTS

In all these patients, EBV serologic tests revealed extremely high IgG antibody titers against early antigen and viral capsid antigen (Table). There were no significant elevations in antibody titers to other herpesviruses or to measles virus in these patients, or to adenovirus in patients 1 and 2.

**Table.** Clinical and laboratory findings in three patients with CEBV

	Patient No.		
	1	2	3
Age (yr)/gender	2/M	6/F	5/F
Prolonged fever	+	+	+
Coronary artery aneurysms	+	+	+
Dilation of sinus of Valsalva	+	-	+
Pericarditis	-	+	+
Rash	-	-	+
Mucous membrane involvement	-	-	-
Desquamation of fingertips	-	-	-
Lymphadenopathy	+	+	-
Hepatosplenomegaly	+	+	+
Liver dysfunction	+	+	+
Interstitial nephritis	-	+	-
Interstitial pneumonitis	+	-	-
Leukopenia	-	-	-
Anemia	+	-	+
Thrombocytopenia	-	-	-
Thrombocytosis	-	-	-
Hypergammaglobulinemia	+	+	+
EBV antibody titer			
VCA-IgG	10,240	20,480	5,120
EA-IgG	2,560	5,120	5,120
EBNA	160	40	10

VCA, Viral capsid antigen; EA, early antigen; EBNA, EBV-determined nuclear antigen.

The PCR-amplified genomic Raji cell DNA was confirmed on ethidium bromide-stained gel and by Southern blot hybridization with a radiolabeled EBV *Bam*HI-W fragment. When DNA samples from cells infected with herpes simplex virus type 1, varicella-zoster virus, human cytomegalovirus, and human herpesvirus-6 were used as

templates in the PCR reaction; no amplification was noted by direct gel analysis or Southern blot hybridization.<sup>6</sup> The EBV genome, as indicated by the 410-base-pair DNA band identical to that of the positive control sample, was detected in cardiac tissue DNA from all three patients and in aortic tissue DNA from one patient (No. 2) on direct gel analysis (Figure). In two control cardiac tissue samples, the PCR procedure did not detect the EBV genome. In tissue samples from the spleen, kidney, and lung of these three patients, the PCR procedure detected the EBV genome. The EBV genome was also detected in aortic tissue of patient 2<sup>7</sup> and in cardiac tissue of patient 3 by Southern blot hybridization without prior DNA amplification.

## DISCUSSION

Vascular and endovascular complications are rare in EBV infection. Loeffel et al.<sup>8</sup> reported an 8-year-old boy with X-linked lymphoproliferative syndrome and aneurysms of central nervous system arteries; the autopsy revealed necrotizing vasculitis in the central nervous system and microscopic changes in the coronary arteries. Ilowite et al.<sup>9</sup> described a 12-year-old boy with Wiskott-Aldrich syndrome who subsequently had pulmonary vasculitis associated with EBV-induced lymphoreticular proliferation; the lung biopsy specimen showed vasculitis involving small to medium-sized arteries and veins.

We now have identified three patients with CEBV accompanied by coronary artery aneurysms. All three had high antibody titers to replicative antigens of EBV and no significant elevations in antibody titers to other herpesviruses, measles virus, or adenovirus. We were not able to test for the presence of the EBV genome in coronary arteries. However, detection of the EBV genome in the heart and aorta suggests that EBV may play a pathogenic role in the development of coronary artery aneurysms in CEBV.

Kawasaki disease is thought to be a self-limited, immunologically mediated vasculitis. We previously reported a link between KD and EBV; 49 (86%) of 57 patients with KD and 15 (68%) of 22 with recurrent KD had serologic evidence of an unusual primary EBV infection.<sup>10,11</sup> The EBV genomes were identified directly by means of the PCR in peripheral blood mononuclear cell DNA samples from 21 (60%) of 35 patients within 2 weeks after the onset of KD. Furthermore, EBV genomes were also detected in all six patients who were repeatedly tested within 3 months after disease onset. In contrast, only 2 (12%) of 17 control DNA samples showed positive PCR results.<sup>6</sup> Epstein-Barr virus genomes were frequently detected in peripheral blood mononuclear cell DNA from KD patients, indicating that there were much higher proportions of EBV-infected cells in their peripheral blood mononuclear cells than in those of

normal control samples. These virologic studies indicate that an unusual EBV-cell interaction may exist in KD.

A broad spectrum of associated features and complications has been reported with CEBV. Coronary artery aneurysms constitute one of the clinical features and appear not to be as rare in these patients as previously assumed. These three patients did not have the clinical hallmarks of KD but all had coronary artery aneurysms. Their recurrent episodes of fever during 6 months to several years can be distinguished from those of atypical KD.<sup>12</sup> All the patients with CEBV associated with coronary artery aneurysms died within 5 years of onset. We recommend periodic echocardiographic examinations for all patients with CEBV. Further studies are needed to explain the relationship between coronary artery aneurysms and EBV infection.

## REFERENCES

1. Schooley RT, Carey RW, Miller G, et al. Chronic Epstein-Barr virus infection associated with fever and interstitial pneumonitis: clinical and serologic features and response to antiviral chemotherapy. *Ann Intern Med* 1986;104:636-43.
2. Straus SE. The chronic mononucleosis syndrome. *J Infect Dis* 1988;157:405-12.
3. Kikuta H, Taguchi Y, Tomizawa K, et al. Epstein-Barr virus genome-positive T lymphocytes in a boy with chronic active EBV infection associated with Kawasaki-like disease. *Nature* 1988;338:455-7.
4. Kawasaki T, Kosaki F, Okawa S, Shigematsu I, Yanagawa H. A new infantile acute mucocutaneous lymph node syndrome (MLNS) prevailing in Japan. *Pediatrics* 1974;54:271-6.
5. Kopolovic J, Pinkus G, Rosen S. Interstitial nephritis in infectious mononucleosis. *Am J Kidney Dis* 1988;12:76-7.
6. Kikuta H, Nakanishi M, Ishikawa N, Konno M, Matsumoto S. Detection of Epstein-Barr virus sequences in patients with Kawasaki disease by means of the polymerase chain reaction. *Intervirology* 1992;33:1-5.
7. Kikuta H, Nakanishi M, Sakiyama Y, Matsumoto S. Chronic active Epstein-Barr virus (EBV) infection is associated with clonotypic intracellular terminal regions of the EBV. *J Infect Dis* 1989;160:546-7.
8. Loeffel S, Chang C-H, Heyn R, et al. Necrotizing lymphoid vasculitis in X-linked lymphoproliferative syndrome. *Arch Pathol Lab Med* 1985;109:546-50.
9. Ilowite NT, Fligner CL, Ochs HD, et al. Pulmonary angitis with atypical lymphoreticular infiltrates in Wiskott-Aldrich syndrome: possible relationship of lymphomatoid granulomatosis and EBV infection. *Clin Immunol Immunopathol* 1986;41:479-84.
10. Kikuta H, Mizuno F, Osato T, et al. Kawasaki disease and an unusual primary infection with Epstein-Barr virus [Letter]. *Pediatrics* 1984;73:413-4.
11. Kikuta H, Matsumoto S, Yanase Y, Kawasaki T, Mizuno F, Osato T. Recurrence of Kawasaki disease and Epstein-Barr virus [Letter]. *J Infect Dis* 1990;162:1215.
12. Rowley AH, Gonzalez-Crussi F, Gidding SS, Duffy CE, Shulman ST. Incomplete Kawasaki disease with coronary artery involvement. *J PEDIATR* 1987;110:409-13.

## Contents JULY 1993

### Original articles

- Development and significance of zidovudine resistance in children infected with human immunodeficiency virus** 1  
*Mark T. Ogino, MD, Wayne M. Dankner, MD, and Stephen A. Spector, MD, San Diego, California*

Resistance to zidovudine was found to be a negative prognostic factor in children infected with human immunodeficiency virus; growth and CD4<sup>+</sup> cell counts were reduced, and death was more likely. Substitution or addition of a second antiretroviral drug was sometimes beneficial. See related article on page 9.

- High-level resistance to zidovudine but not to zalcitabine or didanosine in human immunodeficiency virus from children receiving antiretroviral therapy** 9

*Robert N. Husson, MD, Takuma Shirasaka, MD, PhD, Karina M. Butler, MB, BCh, Philip A. Pizzo, MD, and Hiroaki Mitsuya, MD, PhD, Bethesda, Maryland*

Isolates of human immunodeficiency virus type 1 were evaluated for resistance to antiviral therapy by using a selective polymerase chain reaction technique to identify mutations associated with resistance. Combination therapy did not prevent the emergence of resistance. New methods to prevent the development of resistance to nucleoside analogs are needed. See related article on page 1.

- Prevalence of urinary tract infection in febrile infants** 17  
*Alejandro Hoberman, MD, Han-Pu Chao, MD, David M. Keller, MD, Robert Hickey, MD, Holly W. Davis, MD, and Demetrius Ellis, MD, Pittsburgh, Pennsylvania*

A study of febrile infants seen in an emergency department showed that about 5% had a urinary tract infection, regardless of whether UTI was suspected or other potential causes of fever were identified. The incidence was highest in white girls. Microscopic urinalysis was not a sensitive method of screening; urine culture was required for the diagnosis of UTI.

Vol. 123, No. 1, July 1993. The Journal of Pediatrics (ISSN 0022-3476) is published monthly (six issues per volume, two volumes per year) by Mosby, 11830 Westline Industrial Dr., St. Louis, MO 63146-3318. Second-class postage paid at St. Louis, Mo., and additional mailing offices. POSTMASTER: Send changes of address to The Journal of Pediatrics, Mosby, 11830 Westline Industrial Dr., St. Louis, MO 63146-3318, (800) 453-4351. Annual subscription rates for 1993: domestic, \$88.00 for individuals and \$178.00 for institutions. Printed in the U.S.A. Copyright © 1993 by Mosby-Year Book, Inc. All rights reserved. No part of this publication may be reproduced, stored, or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without permission in writing from the publisher.